

Dr. W. Jean Dodd's vaccination protocol is now being adopted by ALL 27 North American veterinary schools. I highly recommend that you read this. Copy and save it to your files. Print it and pass it out at dog fairs, cat shows, kennel club meetings, dog parks, give a copy to your veterinarian and groomer, etc., etc.* Get the word out.

VACCINATION NEWS FLASH

*I would like to make you aware that all 27 veterinary schools in North America are in the process of changing their protocols for vaccinating dogs and cats. Some of this information will present an ethical & economic challenge to vets, and there will be skeptics.

Some organizations have come up with a political compromise suggesting vaccinations every 3 years to appease those who fear loss of income vs. those concerned about potential side effects. Politics, traditions, or the doctor's economic well being should not be a factor in medical decision.

NEW PRINCIPLES OF IMMUNOLOGY

"Dogs and cats immune systems mature fully at 6 months. If a modified live virus vaccine is given after 6 months of age, it produces an immunity which is good for the life of the pet (ie: canine distemper, parvo, feline distemper). If another MLV vaccine is given a year later, the antibodies from the first vaccine neutralize the antigens of the second vaccine and there is little or no effect. The titer is not "boosted" nor are more memory cells induced." Not only are annual boosters for parvo and distemper unnecessary, they subject the pet to potential risks of allergic reactions and immune-mediated hemolytic anemia. "There is no scientific documentation to back up label claims for annual administration of MLV vaccines." Puppies receive antibodies through their mothers milk. This natural protection can last 8-14 weeks. Puppies & kittens should NOT be vaccinated at LESS than 8 weeks. Maternal immunity will neutralize the vaccine and little protection (0-38%) will be produced. Vaccination at 6 weeks will, however, delay the timing of the first highly effective vaccine. Vaccinations given 2 weeks apart suppress rather than stimulate the immune system. A series of vaccinations is given starting at 8 weeks and given 3-4 weeks apart up to 16 weeks of age. Another vaccination given sometime after 6 months of age (usually at 1 year 4mo) will provide lifetime immunity.

CURRENT RECOMMENDATIONS FOR DOGS

Distemper & Parvo * "According to Dr. Schultz, AVMA, 8-15-95, when a vaccination series given at 2, 3 & 4 months and again at 1 year with a MLV, puppies and kitten program memory cells that survive for life, providing lifelong immunity." Dr. Carmichael at Cornell and Dr. Schultz have studies showing immunity against challenge at 2-10 years for canine distemper & 4 years for parvovirus. Studies for longer duration are pending. "There are no new strains of parvovirus as one mfg. would like to suggest. Parvovirus vaccination provides cross immunity for all types." Hepatitis (Adenovirus) is one of the agents known to be a cause of kennel cough. Only vaccines with CAV-2 should be used as CAV-1 vaccines carry the risk of "hepatitis blue-eye" reactions & kidney damage.

**Bordetella Parainfluenza: Commonly called "Kennel cough" Recommended only for those dogs boarded, groomed, taken to dog shows, or for any reason housed where

exposed to a lot of dogs. The intranasal vaccine provides more complete and more rapid onset of immunity with less chance of reaction. Immunity requires 72 hours and does not protect from every cause of kennel cough. Immunity is of short duration (4 to 6 months).*

*RABIES There have been no reported cases of rabid dogs or cats in Harris, Montgomery or Ft. Bend Counties [Texas], there have been rabid skunks and bats so the potential exists. It is a killed vaccine and must be given every year.

*Lyme disease_is a tick born disease which can cause lameness, kidney failure and heart disease in dogs. Ticks can also transmit the disease to humans. The original Ft. Dodge killed bacteria has proven to be the most effective vaccine. Lyme disease prevention should emphasize early removal of ticks. Amitraz collars are more effective than Top Spot, as amitraz paralyzes the tick's mouthparts preventing transmission of disease.

VACCINATIONS NOT RECOMMENDED

Multiple components in vaccines compete with each other for the immune system and result in lesser immunity for each individual disease as well as increasing the risk of a reaction. Canine Corona Virus is only a disease of puppies. It is rare, self limiting (dogs get well in 3 days without treatment). Cornell & Texas A&M have only diagnosed one case each in the last 7 years. Corona virus does not cause disease in adult dogs.*

*Leptospirosis vaccine is a common cause of adverse reactions in dogs. Most of the clinical cases of lepto reported in dogs in the US are caused by serovars (or types) grippityphosa and bratsilvia. The vaccines contain different serovars eanicola and icthohemorrhagica. Cross protection is not provided and protection is short lived. Lepto vaccine is immuno-suppressive to puppies less than 16 weeks.

NEW DEVELOPMENTS: Giardia is the most common intestinal parasite of humans in North America , 30% or more of all dogs & cats are infected with giardia. It has now been demonstrated that humans can transmit giardia to dogs & cats & vice versa. *

Heartworm preventative must be given year-round in Houston .

*VACCINES BADLY NEEDED

New vaccines in development include: Feline Immunodeficiency Virus and cat scratch fever vaccine for cats and Ehrlichia [one of the other tick diseases, much worse than Lymes] for dogs.

THE VIEW FROM THE TRENCHES; BUSINESS ASPECTS

Most vets recommend annual boosters and most kennel operators require them. For years the pricing structure of vets has misled clients into thinking that the inherent value of an annual office visit was in the "shots" they failed to emphasize the importance of a physical exam for early detection of treatable diseases. It is my hope that you will continue to require rabies & Kennel cough and emphasize the importance of a recent vet exam. I also hope you will accept the new protocols and honor these pets as currently vaccinated. Those in the boarding business who will

honor the new vaccine protocols can gain new customers who were turned away from vet owned boarding facilities reluctant to change.

CONCLUSION

Dogs & cats no longer need to be vaccinated against distemper, parvo, & feline leukemia every year. Once the initial series of puppy or kitten vaccinations and first annual vaccinations are completed, immunity from MLV vaccines persists for life. It has been shown that cats over 1 year of age are immune to Feline Leukemia whether they have been vaccinated or not. Imagine the money you will save, not to mention less risks from side effects. PCR rabies vaccine, because it is not adjuvanted, will mean less risk of mediated hemolytic anemia and allergic reactions are reduced by less frequent use of vaccines as well as by avoiding unnecessary vaccines such as K-9 Corona virus and chlamydia for cats, as well as ineffective vaccines such as Leptospirosis and FIP. Intranasal vaccine for Rhotracheitis and Calici virus, two upper respiratory viruses of cats provide more complete protection than injectable vaccines with less risk of serious reactions.

The AAHA and all 27 veterinary schools of North America are our biggest endorsement for these new protocols.*

Dr. Bob Rogers

Please consider as current on all vaccinations for boarding purposes.

DOGS Initial series of puppy vaccines

1. distemper, hepatitis, parvo, parinfluenza - 3 sets one month apart concluding at 16 weeks of age.*

2. Rabies at 16 weeks of age (later is better)

3. Bordetella within last 4-6 months First annual (usually at 1 year and 4 months of age)*

1. DHP, Parvo, Rabies

2. Bordetella within last 4-6 months 2 years or older

1. Rabies with in last year

2. Bordetella within last 4-6 months

3. DHP & Parvo given anytime over 6 months of age , but not necessarily within the last year.

Recommended: Physical exam for transmissible diseases and health risks.